



Guidance on Infection Control In Schools and other Child Care Settings

Prevent the spread of infections by ensuring: routine immunisation, high standards of personal hygiene and practice, particularly hand washing, and maintaining a clean environment.

Diarrhoea and Vomiting illness#

	Recommended period to be kept away from school, nursery, or childminders	Comments
Diarrhoea and/or vomiting	48 hours from last episode of diarrhoea or vomiting (48hr rule applies).	Exclusion from swimming should be for 2 weeks following last episode of diarrhoea.
E. coli 0157 VTEC	Exclusion is important for some children. Always consult with HPU.	Exclusion applies to young children and those who may find hygiene practices difficult to adhere to. Local HPU will advise. Exclusion from swimming should be for 2 weeks following last episode of diarrhoea.
Typhoid* [and paratyphoid*] (enteric fever)	Exclusion is important for some children. Always consult with HPU.	Exclusion applies to young children and those who may find hygiene practices difficult to adhere to. Local HPU will advise. Exclusion from swimming should be for 2 weeks following last episode of diarrhoea.
Shigella (Dysentery)	Exclusion may be necessary.	Exclusion (if required) applies to young children and those who may find hygiene practices difficult to adhere to. Local HPU will advise. Exclusion from swimming should be for 2 weeks following last episode of diarrhoea.

Respiratory Infections

'Flu' (influenza)	Until recovered.	SEE: vulnerable children.
Tuberculosis*	Always consult with HPU.	Not usually spread from children. Requires quite prolonged, close contact for spread.
Whooping cough* (Pertussis)	Five days from commencing antibiotic treatment or 21 days from onset of illness if no antibiotic treatment.	Preventable by vaccination. After treatment non-infectious coughing may continue for many weeks. HPU will organise any contact tracing necessary.